

ACB Yerba Santa Glycoprotein PF Efficacy Data

Code: 20342PF
INCI Name: Lactobacillus/Eriodictyon Californicum Ferment Filtrate
CAS #: 68990-14-7
EINECS #: 273-580-8

Name of Study	Results
Moisturization Study	The chart shows the results of the % moisturization at day 1 and day 30 of the study. The graph shows the results of the percent increase in moisturization comparing Aloe Vera and Yerba Santa Glycoprotein.
Cellular Viability Assay	In this study, ACB Yerba Santa Glycoprotein PF (code 20342PF) was tested to evaluate its effects on the viability of normal human dermal fibroblasts (NDHF). At concentrations of both 0.1% and 0.01% ACB Yerba Santa Glycoprotein PF (code 20342PF), nor the preservatives contained therein exhibited any inhibition of cell viability. It can therefore be concluded that at normal use concentrations ACB Yerba Santa Glycoprotein PF (code 20342PF) is not cytotoxic.
ORAC Assay	As shown in figure 1, ACB Yerba Santa Glycoprotein PF exhibited antioxidant activity comparable to 200µM Trolox®. The antioxidant capacity of ACB Yerba Santa Glycoprotein PF increased as the concentration increased. As a result we can assure that its ability to minimize oxidative stress is dose dependent. With the present study we can confirm that this unique ingredient is capable of providing antioxidant benefits when added to cosmetic applications.



Moisturization/Hydration Assay

info@activeconceptsllc.com • Phone: +1-704-276-7100 • Fax: +1-704-276-7101

Tradename: ACB Yerba Santa Glycoprotein PF

Code: 20342PF

CAS #: 68990-14-7

Test Request Form #: 2782

Lot #: 32061

Sponsor: *Active Concepts, LLC; 107 Technology Drive Lincolnton, NC 28092*

Study Director: *Maureen Danaher*

Principle Investigator: *Jennifer Goodman*

Introduction:

An *in-vivo* clinical Twelve- (12) subject study was carried out over a thirty-day period to evaluate the long-term effects of the cosmetic raw material, **ACB Yerba Santa Glycoprotein PF** on skin moisturization.

Materials:

Product 1: Blank Formulation Base (Water) + Aloe Vera 10X Gel (Control)

Product 2: Blank Formulation Base (Water) + 5% **ACB Yerba Santa Glycoprotein PF**

Equipment:

Nova Impedance Meter

Methods:

Active Concepts utilized 12 female volunteers between the ages of 18 and 45, known to be free of any skin pathologies. On day one of the study the moisturization levels of the volunteers' skin were measured via Nova Impedance Meter. This measurement was used as the subject's baseline. We then divided the volunteers into two separate groups – Group 1 (6 volunteers) used Product 1 on the right side of the face. The left side of the face remained untreated. Group 2 (6 volunteers) used Product 2 on the right side of the face. The left side of the face remained untreated. The twelve subjects followed the protocol for thirty days, applying the products once daily on the appropriate side of their face. On day 30 Moisturization was once again assessed on the two halves of the face of each of the volunteers using the Nova Impedance Meter. The amount of moisturization obtained after the 30 day study was evaluated and compared to the measurements taken on day one the study. Significant differences were found between the two groups of volunteers.


This information is presented in good faith but is not warranted as to accuracy of results. Also, freedom from patent infringement is not implied.
This information is offered solely for your investigation, verification, and consideration.

Results/Conclusion:

The following chart shows the results of the % moisturization at day 1 and day 30 of the study. The graph shows the results of the percent increase in moisturization comparing Aloe Vera and **ACB Yerba Santa Glycoprotein PF**.

Group 1(Aloe)	
Day 1 Average % Moisturization	5.20%
Day 30 Average % Moisturization	28%
Group 2(ACB Yerba Santa Glycoprotein PF)	
Day 1 Average % Moisturization	5.40%
Day 30 Average % Moisturization	37%

Chart 1: Percent moisturization.



Graph 1: Total increase in moisturization.



Cellular Viability Assay Analysis

info@activeconceptsllc.com • Phone: +1-704-276-7100 • Fax: +1-704-276-7101

Tradename: ACB Yerba Santa Glycoprotein PF

Code: 20342PF

CAS #: 68990-14-7

Test Request Form #: 622

Lot #: 22061

Sponsor: *Active Concepts, LLC; 107 Technology Drive Lincolnton, NC 28092*

Study Director: *Erica Segura*

Principle Investigator: *Meghan Darley*

Test Performed:

Cellular Viability Assay

Introduction

The cellular viability assay is useful for quantitatively measuring cell-mediated cytotoxicity, cell proliferation and mitochondrial metabolic activity. Increased metabolism in a cell indicates ample cellular respiration and adenosine triphosphate (ATP) production. ATP is the molecular energy of cells and is required in basic cell function and signal transduction. A decrease in ATP levels indicates cytotoxicity and decreased cell function while an increase in ATP levels indicates healthy cells.

The cellular viability assay was conducted to assess the ability of **ACB Yerba Santa Glycoprotein PF** to increase cellular metabolic activity in cultured dermal fibroblasts.

Assay Principle

The assay utilizes a nonfluorescent dye, resazurin, which is converted to a fluorescent dye, resorufin, in response to chemical reduction of growth medium from cell growth and by respiring mitochondria. Healthy cells that are in a proliferative state will be able to easily convert resazurin into resorufin without harming the cells. This method is a more sensitive assay than other commonly used mitochondrial reductase dyes such as MTT. An increase in the signal generated by resazurin-conversion is indicative of a proliferative cellular state.

This information is presented in good faith but is not warranted as to accuracy of results. Also, freedom from patent infringement is not implied.
This information is offered solely for your investigation, verification, and consideration.



Cellular Viability Assay Analysis

info@activeconceptsllc.com • Phone: +1-704-276-7100 • Fax: +1-704-276-7101

Materials

- A. **Kit:** PrestoBlue™ Cell Viability Reagent (Invitrogen, A13261)
- B. **Incubation Conditions:** 37°C at 5% CO₂ and 95% relative humidity (RH)
- C. **Equipment:** Forma humidified incubator; ESCO biosafety laminar flow hood; Light microscope; Pipettes
- D. **Cell Line:** Normal Human Dermal Fibroblasts (NHDF) (Lonza; CC-2511)
- E. **Media/Buffers:** Dulbecco's Modified Eagle Medium (DMEM); Penicillin-Streptomycin (50U-50mg/mL); Fetal Bovine Serum (FBS); Phosphate Buffered Saline (PBS)
- F. **Culture Plate:** Falcon flat bottom 96-well tissue culture treated plates
- G. **Reagents:** PrestoBlue™ reagent (10X)
- H. **Other:** Sterile disposable pipette tips

Methods

Human dermal fibroblasts were seeded into 96-well tissue culture plates and allowed to grow to confluency in complete DMEM. A 10-fold serial dilution was performed resulting in **ACB Yerba Santa Glycoprotein PF** concentrations of 0.1% and 0.01% in complete DMEM and incubated with fibroblasts for 24 hours.

Ten microliters of viability reagent was added to 90µL of cell culture media in culture wells and a fluorometric measurement was taken at 560nm for excitation and 590nm for emission.

This information is presented in good faith but is not warranted as to accuracy of results. Also, freedom from patent infringement is not implied. This information is offered solely for your investigation, verification, and consideration.

Results

The data obtained from this study met criteria for a valid assay and the controls performed as anticipated.

ACB Yerba Santa Glycoprotein PF did not exhibit significant effects on the cellular metabolism.

Cellular metabolism results are shown as mean fluorescence units (MFU) and expressed as percentage change, calculated by the below equation:

$$\text{Percent (\%)Change} = \frac{MFU_{Control} - MFU_{Sample}}{MFU_{Control}} \times 100$$




Figure 1: Cellular Metabolism of **ACB Yerba Santa Glycoprotein PF**-treated fibroblasts expressed in terms of percent of control.

Discussion

In this study, **ACB Yerba Santa Glycoprotein PF** (code 20342PF) was tested to evaluate its effects on the viability of normal human dermal fibroblasts (NDHF). At concentrations of both 0.1% and 0.01% **ACB Yerba Santa Glycoprotein PF** (code 20342PF), nor the preservatives contained therein exhibited any inhibition of cell viability. It can therefore be concluded that at normal use concentrations **ACB Yerba Santa Glycoprotein PF** (code 20342PF) is not cytotoxic.



Oxygen Radical Absorbance Capacity (ORAC) Assay

info@activeconceptsllc.com • Phone: +1-704-276-7100 • Fax: +1-704-276-7101

Tradename: ACB Yerba Santa Glycoprotein PF

Code: 20342PF

CAS #: 68990-14-7

Test Request Form #: 596

Lot #: 32061

Sponsor: *Active Concepts, LLC; 107 Technology Drive Lincolnton, NC 28092*

Study Director: *Erica Segura*

Principle Investigator: *Meghan Darley*

Test Performed:

Oxygen Radical Absorbance Capacity (ORAC)

Introduction

Reactive oxygen species (ROS) are generated by normal cellular processes, environmental stresses, and UV irradiation. ROS are detrimental to cellular structures and functional molecules (i.e DNA, proteins, lipids) as they act as strong oxidizing agents or free radicals. The oxygen radical absorbance capacity (ORAC) assay is a standard method used to assess antioxidant capacity of physiological fluids, foods, beverages, and natural products. The assay quantitatively measures a sample's ability to quench free radicals that have the potential to react with and damage cellular components.

Oxygen Radical Absorbance Capacity (ORAC) assay was conducted to assess the antioxidant capacity of **ACB Yerba Santa Glycoprotein PF**.

Assay Principle

This assay is based upon the effect of peroxy radicals generated from the thermal decomposition of 2, 2'-azobis-2-methyl-propanimidamide dihydrochloride (AAPH) on the signal intensity from the fluorescent probe, fluorescein, in the presence of an oxygen radical absorbing substance. The degree of change is indicative of the amount of radical damage and the presence of antioxidants results in an inhibition in the free radical damage to the fluorescein. The antioxidant protection of the sample can be calculated by comparing it to a set of known standards. Trolox®, a water soluble vitamin E analog, with known antioxidant capabilities is used in this ORAC assay as the standard for measuring the antioxidant capacity of unknown substances. ORAC values, expressed in μM of Trolox® equivalents (TE), are calculated using the area under the curves (AUC) of the test product, Trolox®, and the control materials. Trolox equivalency is used as the benchmark for antioxidant capacity of mixtures since it is difficult to measure individual components.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.



Oxygen Radical Absorbance Capacity (ORAC) Assay

info@activeconceptsllc.com • Phone: +1-704-276-7100 • Fax: +1-704-276-7101

Materials

- A. Equipment:** Synergy H1 Microplate reader (BioTek Instruments, Winooski, VT); Gen5 software (BioTek Instruments, Winooski, VT); Pipettes
- B. Buffers:** 75mM Potassium Phosphate (pH 7.4); Deionized H₂O
- C. Reagents:** 2,2'-Azobis(2-methylpropionamide) dihydrochloride (AAPH) (153mM); 6-Hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid (Trolox®); Fluorescein Sodium Salt (4nM)
- D. Preparation:** Pre-heat (37°C) Synergy H1 Microplate reader; Prepare Trolox® standards, sample dilutions, fluorescein solution, and AAPH.
- E. Microtitre Plates:** Corning 96 Well Black Side/Clear Bottom Microplates

Methods

Solutions of **ACB Yerba Santa Glycoprotein PF** and Trolox® (positive control) were prepared in 75mM potassium phosphate buffer. Materials were prepared at three different concentrations/dilutions. Trolox® was used as a reference for antioxidant capacity and prepared at a concentrations ranging from 12.5µM to 200µM in 75mM potassium phosphate buffer.

For the ORAC assay, 25µL of test material and Trolox® were combined with 150µL of fluorescein in 75mM potassium phosphate buffer and incubated in the Synergy HT Microplate reader at 37°C for 30 minutes. At the end of the incubation period, 25µL of AAPH were pipetted into each well. Fluorescent measurements were then taken every 2 minutes for approximately 2 hours at an excitation wavelength of 485nm and an emissions wavelength of 520nm.

The AUC and Net AUC values of the standards and samples were determined using Gen5 2.0 Data Reduction Software using the below equations:

$$AUC = 0.5 + \frac{R2}{R1} + \frac{R3}{R1} + \frac{R4}{R1} + \dots + \frac{Rn}{R1} \rightarrow \text{Where } R \text{ is fluorescence reading}$$

$$\text{Net AUC} = AUC_{\text{sample}} - AUC_{\text{blank}}$$

The standard curve was obtained by plotting the Net AUC of different Trolox® concentrations against their concentration. ORAC values of samples were then calculated automatically using the Gen5 software to interpolate the sample's Net AUC values against the Trolox® standard curve. ORAC measurements for the test material were expressed in micro moles Trolox® equivalents (µMTE), where 1 ORAC unit is equal to 1 µMTE.

ORAC values are also calculated in Units/milliliter (U/mL). The equation used for the calculation is shown below:

$$\text{ORAC (U/mL)} = (50 \times \text{Dilution Factor}) \times \left(\frac{AUC_{\text{Sample}} - AUC_{\text{Blank}}}{AUC_{\text{Trolox}} - AUC_{\text{Blank}}} \right)$$

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.



Oxygen Radical Absorbance Capacity (ORAC) Assay

info@activeconceptsllc.com • Phone: +1-704-276-7100 • Fax: +1-704-276-7101

Results

ACB Yerba Santa Glycoprotein PF began exhibiting antioxidant activity at a 0.05% concentration.

At concentrations exceeding 0.1%, the antioxidant activity is too intense to measure, resulting in over-saturation of the signal. This over-saturation produces results that exceed those of the standard curve. Therefore, lower concentrations were tested to ensure an accurate result.

The ORAC value expressed in U/mL for 0.1% **ACB Agave HSP** is 39382.9.




Figure 1: Antioxidant capacities

Discussion

As shown in figure 1, **ACB Yerba Santa Glycoprotein PF** exhibited antioxidant activity comparable to 200µM Trolox®. The antioxidant capacity of **ACB Yerba Santa Glycoprotein PF** increased as the concentration increased. As a result we can assure that its ability to minimize oxidative stress is dose dependent. With the present study we can confirm that this unique ingredient is capable of providing antioxidant benefits when added to cosmetic applications.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.