

**Tradename:** Probacillus Revive

**Code:** 16618

**CAS #:** 9015-54-7

**Test Request Form #:** 6014

**Lot #:** 75876

**Sponsor:** *Active Concepts, LLC; 107 Technology Drive Lincolnton, NC 28092*

**Study Director:** *Maureen Danaher*

**Principal Investigator:** *Michael Hovis*

**Test Performed:**

Scratch Assay

## **Introduction**

Wounded tissue begins a complex and structured series of events in order to repair the damaged region. Some of these events include upregulation of angiogenic factors causing increased vascularization, increased deposition of extracellular matrix, and increased cell proliferation. The wound healing process begins as cells polarize toward the wound, initiate protrusion, migrate, and close the wound area. These processes reflect the behavior of individual cells as well as the entire tissue complex.

A Scratch Assay was conducted to assess the *in vitro* wound healing properties of **Probacillus Revive** in dermal fibroblasts.

## **Assay Principle**

The *in vitro* scratch assay is a well-known and widely used method to study cell migration and proliferation. This assay is based on the observation that when an artificial gap or scratch is made on a confluent cell monolayer, the cells will migrate towards the opening and close the scratch. The basic steps involve creating a scratch in a cell monolayer and capturing images throughout the healing or cell migration process. Through these images we can quantify the rate of cell migration.

## Materials

- A. Incubation Conditions:** 37°C, 5% CO<sub>2</sub>, and 95% relative humidity (RH)
- B. Equipment:** Forma Humidified Incubator; ESCO Biosafety Laminar Flow Hood; Camera; Pipettes; Light Microscope
- C. Cell Line:** Normal Human Dermal Fibroblasts (ATCC; PCS-201-012)\*
- D. Media/Buffers:** Fibroblast Basal Medium (PCS-201-030)\*; Fibroblast Growth Kit (PCS-201-041)\*; Phosphate Buffered Saline (PBS)
- E. Reagents:** Epidermal Growth Factor-1 (100 ng/mL); Paraformaldehyde (3.7%); Crystal Violet Stain
- F. Culture Plate:** 6 Well Flat Bottom Tissue Culture Treated Plates
- G. Analysis Software:** ImageJ (National Institutes of Health); Excel Analysis ToolPak (Microsoft)
- H. Other:** Sterile disposable pipette tips
- \*Or suitable alternatives, subject to change without notice based off vendor availability

## Methods

Human dermal fibroblasts were seeded into a 6-well tissue culture plate and allowed to grow to confluency in complete media. When cell growth reached confluency, scratches were made down the middle of the well in a straight line, generating an in vitro “wound” which is the area devoid of cells. The rate at which the fibroblasts migrate to fill the area devoid of cells indicates wound healing. All wells were washed with sterile PBS to remove cellular debris caused by the scratch. The positive control, Epidermal Growth Factor-1 (EGF-1) (100 ng/mL), and a market standard healing active, Vitamin E (0.02% and 0.05%), were diluted with serum-free complete media. A 0.1% concentration of **Probacillus Revive** were diluted with serum-free complete media. Serum-free complete media was used as a negative control. Media was not changed throughout the duration of the experiment. Images were captured immediately after the scratch (t=0) and every 24-hours afterwards, up to 72 hours. Cells were fixed with 3.7% paraformaldehyde and stained with crystal violet for enhanced microscopy.

ImageJ software was used to analyze the images and calculate the area of the scratch and the closure rate. Percent scratch closure and migration rate were calculated by the following formulas:

$$\frac{\text{Scratch Area}_{t=x} - \text{Scratch Area}_{t=0}}{\text{Scratch Area}_{t=0}} \times 100 = \% \text{ Scratch Closure}$$

$$\frac{\text{Change in Area of Scratch (nm}^2\text{)}}{\text{Migration Time}_{t=x}} = \text{Migration Rate}$$

$$x = \text{time (hours) post scratch}$$


Area under the curve was calculated for percent scratch closure with the following equation:

$$AUC = \frac{1}{2} \sum_{i=0}^{n-1} (t_{i+1} - t_i) (\text{Percent Closure}_i + \text{Percent Closure}_{i+1})$$


For percent scratch closure calculations *Percent Closure*<sub>1</sub> and *Percent Closure*<sub>2</sub> at times *t*<sub>1</sub> and *t*<sub>2</sub>, the AUC between those two time points is equivalent to the product of difference in time and the average of the two percent scratch closure calculations. Three separate experiments were performed with conditions in duplicate and average values were recorded. Data was analyzed using a one-way ANOVA with statistical significance accepted at *p* ≤ 0.05.

## Results

The data obtained from this study met criteria for a valid assay and the positive and negative controls performed as anticipated. Fibroblasts treated with 0.1% **Probacillus Revive** increased cell migration and wound healing compared to the negative control (serum-free complete media).



**Figure 1.** Effect of **Probacillus Revive** on the Area of Scratch over time.



**Figure 2.** Effect of **Probacillus Revive** on the Percent of Scratch Closure over time.




Figure 3. Effect of Probacillus Revive on the Rate of Migration.





Figure 4. Images of EGF-1 (positive control), Serum Free Complete Media (negative control), and 0.1% Probacillus Revive at T=0 (A, D, G), T=72 (B, E, H), and stained at T=72 (C, F, I).



**Figure 5.** Area Under the Curve (AUC) of Percent Scratch Closure After 72 Hours of Treatment. \*indicates significance ( $p \leq 0.05$ ) compared to Serum Free Media. ^ indicates significance ( $p \leq 0.05$ ) compared to Complete Media.

**Table 1.** Results from one-way ANOVA Statistical Analysis of Total Percent Scratch Closure (Percent AUC). Results represent p-values between the two conditions compared. \* indicates significance ( $p \leq 0.05$ ) compared to Serum Free Media. ^ indicates significance ( $p \leq 0.05$ ) compared to Complete Media.

	Serum-Free Complete Media	Complete Media	EGF-1 (100 ng/mL)	0.1% Probacillus Revive
Serum-Free Complete Media	-----	0.001*^	< 0.001*	< 0.001*
Complete Media	0.001*^	-----	0.038^	0.041^
EGF-1 (100 ng/mL)	< 0.001*	0.038^	-----	> 0.05
0.1% Probacillus Revive	< 0.001*	0.041^	> 0.05	-----

## Discussion

As shown in Figures 1-5, fibroblasts incubated with EGF-1, a molecule known to augment cellular migration, enhanced wound healing compared to untreated fibroblasts. Conversely, fibroblasts exposed to serum-free complete media, the negative control, elicited a reduction in cell migration compared to untreated fibroblasts. These data demonstrate cellular migration in fibroblasts is dynamic and can be manipulated with exogenous substances.

Similarly, fibroblasts treated with 0.1% **Probacillus Revive** increased cell migration and closed the scratch at a rate comparable to the positive control (EGF-1) and greater than untreated fibroblasts (Table 1). These data demonstrate **Probacillus Revive** activates cellular migration and wound healing in fibroblasts.

Collectively, the mechanisms of *in vitro* scratch closure mimic the mechanisms seen in *in vivo* wound healing. These data indicate **Probacillus Revive** has wound healing properties and triggers cellular migration, which may assist in the wound healing process.